



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

$$+ \cot^2 \frac{\pi}{2n} \log \left( \frac{1 + \sin \frac{\pi}{2n}}{\cos \frac{\pi}{2n}} \right) \Big].$$

COROLLARY. If  $a=1$ ,  $n=2$ , then

$$M = \frac{2 - \sqrt{2}}{3} + \log(1 + \sqrt{2}).$$

Also solved by *G. B. M. ZERE*, and *HENRY HEATON*.

### MISCELLANEOUS.

90. Proposed by *DR. E. D. ROE, Jr.*, Syracuse University, Syracuse, N. Y.

I shot my rifle at different ranges and found the following table of elevations  $e$ , for the vernier peep sight, for the given distances  $s$ :

$s$	$e$
0	21.0
100	24.5
200	28.5
300	33.5
400	40.0
500	48.5

The distances are measured in yards. How shall a table of elevations be constructed, giving the arguments  $e$ , for every five yards up to 500 yards? Do not give the whole table, but explain the method, and illustrate by giving a computation, carrying the result to three places of decimals. An actual problem.

Solution by the **PROPOSER**.

The solution of this problem, which is doubtless somewhat arbitrary, was as follows: We have partly given and partly implied in the data, the following scheme:

$s$	0	100	200	300	400	500
$e$	21.0	24.5	28.5	33.5	40.0	48.5
$\Delta e$	3.5	4.0	5.0	6.5	8.5	
$\Delta^2 e$	0.5	1.0	1.5	2.0		
$\Delta^3 e$	0.5	0.5	0.5			
$\Delta^4 e$	0	0				

In this we notice that the fourth and third differences are constant, while the second are in arithmetical progression. Assuming that we have discovered the law, we may extend the second, third and fourth differences up to the column under 400, and thus have the complete data for interpolation by means of the method of finite differences. According to this method, if  $e_s$  denote the elevation for the distance  $s$ , we have,

$$e_y \times 100 + (x/20) \times 100 = e_{100y+5x} = e_{100y} + \frac{x}{20} \frac{\Delta e}{1!} 100y + \frac{x}{20} \left( \frac{x}{20} - 1 \right) \frac{\Delta^2 e}{2!} 100y \\ + \frac{x}{20} \left( \frac{x}{20} - 1 \right) \left( \frac{x}{20} - 2 \right) \frac{\Delta^3 e_{100y}}{3!},$$

where  $y=0, 1, 2, 3, 4$ .

$x=1, 2, \dots, 19$ . As the complete solution, and giving the 95 values sought.

Example: It is required to find  $e_{425}$ . Here  $y=4$ ,  $x=5$ , and  $e_{400}=40$ ,  $\Delta e_{400}=8.5$ ,  $\Delta^2 e_{400}=2.5$ ,  $\Delta^3 e_{400}=0.5$ ,  $\Delta^4 e_{400}=0$ . Hence

$$e_{425} = 40 + \frac{1}{4} 8.5 + \frac{1}{4} \left( \frac{1}{4} - 1 \right) \frac{2.5}{2!} + \frac{1}{4} \left( \frac{1}{4} - 1 \right) \left( \frac{1}{4} - 2 \right) \frac{0.5}{3!} \\ = 40 + 2.125 - 0.234 + 0.027 = 41.918.$$

Remark: With a table like this satisfactory results in hunting may be obtained. The table is contained in a small note book. But the style of hunting must be changed. Longer distances must be used, and the work resolves itself into judging distances, and variations in the wind, setting elevation and wind gauge sights, accurate sighting and firm arm holding of the rifle.

Also solved by *G. B. M. ZERR*.

---

## PROBLEMS FOR SOLUTION.

---

### ARITHMETIC.

144. Proposed by *B. F. FINKEL*, A. M., M. Sc., Professor of Mathematics and Physics in Drury College, Springfield, Mo.

A hired a house for one year for \$300; at the end of four months he takes in M as a partner; and at the end of eight months he takes in P. At the end of the year what rent must each pay? [From Greenleaf's *National Arithmetic*, page 442.]

145. Proposed by *B. F. FINKEL*, A. M., M. Sc., Professor of Mathematics and Physics in Drury College, Springfield, Mo.

By discounting a note at 20% per annum, I get  $22\frac{1}{2}\%$  per annum interest; how long does the note run? [From Ray's *Higher Arithmetic*, page 405.]

\*\*\* Solutions of these problems should be sent to B. F. Finkel not later than July 10.

---

### ALGEBRA.

136. Proposed by *JOHN M. COLAW*, A. M., Monterey, Va.

Solve  $a^x b^y = c \dots (1)$ , and  $c^{x+y} = ab \dots (2)$ .

137. Proposed by *MARCUS BAKER*, U. S. Geological Survey, Washington, D. C.

Solve, if possible,  $a^x + b^x = c$ .